

## A NOTE ON ADDITIVITY OF POLYGAMMA FUNCTIONS

FENG QI AND BAI-NI GUO

ABSTRACT. In the note, the functions  $|\psi^{(i)}(e^x)|$  for  $i \in \mathbb{N}$  are proved to be sub-additive on  $(\ln \theta_i, \infty)$  and super-additive on  $(-\infty, \ln \theta_i)$ , where  $\theta_i \in (0, 1)$  is the unique root of equation  $2|\psi^{(i)}(\theta)| = |\psi^{(i)}(\theta^2)|$ .

## 1. INTRODUCTION

Recall [3, 5, 7] that a function  $f$  is said to be sub-additive on  $I$  if

$$f(x+y) \leq f(x) + f(y) \quad (1)$$

holds for all  $x, y \in I$  such that  $x+y \in I$ . If the inequality (1) is reversed, then  $f$  is called super-additive on  $I$ .

The sub-additive and super-additive functions play an important role in the theory of differential equations, in the study of semi-groups, in number theory, and also in the theory of convex bodies. A lot of literature for the sub-additive and super-additive functions can be found in [3, 5] and related references therein.

It is well-known that the classical Euler gamma function  $\Gamma(x)$  may be defined for  $x > 0$  by

$$\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt. \quad (2)$$

The logarithmic derivative of  $\Gamma(x)$ , denoted by  $\psi(x) = \frac{\Gamma'(x)}{\Gamma(x)}$ , is called the psi or digamma function, and  $\psi^{(k)}(x)$  for  $k \in \mathbb{N}$  are called the polygamma functions. It is common knowledge that these functions are fundamental and important and that they have much extensive applications in mathematical sciences.

In [4], the function  $\psi(a+x)$  is proved to be sub-multiplicative with respect to  $x \in [0, \infty)$  if and only if  $a \geq a_0$ , where  $a_0$  denotes the only positive real number which satisfies  $\psi(a_0) = 1$ .

In [5], the function  $[\Gamma(x)]^\alpha$  was proved to be sub-additive on  $(0, \infty)$  if and only if  $\frac{\ln 2}{\ln \Delta} \leq \alpha \leq 0$ , where  $\Delta = \min_{x \geq 0} \frac{\Gamma(2x)}{\Gamma(x)}$ .

In [2, Lemma 2.4], the function  $\psi(e^x)$  was proved to be strictly concave on  $\mathbb{R}$ .

In [7, Theorem 3.1], the function  $\psi(a+e^x)$  is proved to be sub-additive on  $(-\infty, \infty)$  if and only if  $a \geq c_0$ , where  $c_0$  is the only positive zero of  $\psi(x)$ .

In [6, Theorem 1], among other things, it was presented that the function  $\psi^{(k)}(e^x)$  for  $k \in \mathbb{N}$  is concave (or convex, respectively) on  $\mathbb{R}$  if  $k = 2n - 2$  (or  $k = 2n - 1$ , respectively) for  $n \in \mathbb{N}$ .

In this short note, we discuss sub-additive and super-additive properties of polygamma functions  $\psi^{(i)}(x)$  for  $i \in \mathbb{N}$ .

---

2000 *Mathematics Subject Classification.* Primary 33B15; Secondary 26D07.

*Key words and phrases.* sub-additive function, super-additive function, polygamma function.

This paper was typeset using  $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$ .

Our main result is the following Theorem 1.

**Theorem 1.** *The functions  $|\psi^{(i)}(e^x)|$  for  $i \in \mathbb{N}$  are super-additive on  $(-\infty, \ln \theta_i)$  or sub-additive on  $(\ln \theta_i, \infty)$ , where  $\theta_i \in (0, 1)$  is the unique root of equation*

$$2|\psi^{(i)}(\theta)| = |\psi^{(i)}(\theta^2)|. \quad (3)$$

## 2. PROOF OF THEOREM 1

Let

$$f(x, y) = |\psi^{(i)}(x)| + |\psi^{(i)}(y)| - |\psi^{(i)}(xy)| \quad (4)$$

for  $x > 0$  and  $y > 0$ , where  $i \in \mathbb{N}$ . In order to show Theorem 1, it is sufficient to prove the positivity or negativity of the function  $f(x, y)$ . Direct differentiation yields

$$\begin{aligned} \frac{\partial f(x, y)}{\partial x} &= y|\psi^{(i+1)}(xy)| - |\psi^{(i+1)}(x)| \\ &= \frac{1}{x} [xy|\psi^{(i+1)}(xy)| - x|\psi^{(i+1)}(x)|]. \end{aligned} \quad (5)$$

In [1, Lemma 1] and [8, 9], among other things, the functions  $x^\alpha |\psi^{(i)}(x)|$  are proved to be strictly increasing on  $(0, \infty)$  if and only if  $\alpha \geq i + 1$  and strictly decreasing if and only if  $\alpha \leq i$ . From this monotonicity, it follows easily that  $\frac{\partial f(x, y)}{\partial x} \geq 0$  if and only if  $y \leq 1$ , which means that the function  $f(x, y)$  is strictly increasing for  $y < 1$  and strictly decreasing for  $y > 1$  in  $x \in (0, \infty)$ . Since

$$\lim_{x \rightarrow \infty} f(x, y) = |\psi^{(i)}(y)| > 0,$$

then the function  $f(x, y)$  is positive in  $x \in (0, \infty)$  for  $y > 1$ .

For  $y < 1$ , by virtue of the increasing monotonicity of  $f(x, y)$ , it is deduced that

- (1) if  $x > 1$ , then  $f(1, y) = |\psi^{(i)}(1)| < f(x, y) < |\psi^{(i)}(y)|$ ;
- (2) if  $x < 1$ , then  $f(x, y) < f(1, y) = |\psi^{(i)}(1)|$ ;
- (3) if  $y < x < 1$ , then  $f(y, y) < f(x, y)$ ;
- (4) if  $x < y < 1$ , then  $f(x, x) < f(x, y)$ .

This implies that

$$f(\theta, \theta) = 2|\psi^{(i)}(\theta)| - |\psi^{(i)}(\theta^2)| < f(x, y) \quad (6)$$

for  $y < 1$ , where  $\theta < 1$  with  $\theta < x$  and  $\theta < y$ . Since  $f(\theta, \theta)$  is strictly increasing on  $(0, 1)$  such that  $f(1, 1) = |\psi^{(i)}(1)| > 0$  and  $\lim_{\theta \rightarrow 0^+} f(\theta, \theta) = -\infty$ , then the function  $f(\theta, \theta)$  has a unique zero  $\theta_i \in (0, 1)$  such that  $f(\theta, \theta) > 0$  for  $1 > \theta > \theta_i$ .

In conclusion, the function  $f(x, y)$  is positive for  $x, y > \theta_i$  or negative for  $0 < x, y < \theta_i$ . The proof of Theorem 1 is complete.

## REFERENCES

- [1] H. Alzer, *Mean-value inequalities for the polygamma functions*, Aequationes Math. **61** (2001), no. 1, 151–161.
- [2] H. Alzer, *Sharp inequalities for the digamma and polygamma functions*, Forum Math. **16** (2004), 181–221.
- [3] H. Alzer and S. Koumandos, *Sub- and super-additive properties of Fejér's sine polynomial*, Bull. London Math. Soc. **38** (2006), no. 2, 261–268.
- [4] H. Alzer and O. G. Ruehr, *A submultiplicative property of the psi function*, J. Comput. Appl. Math. **101** (1999), 53–60.

- [5] H. Alzer and S. Ruscheweyh, *A subadditive property of the gamma function*, J. Math. Anal. Appl. **285** (2003), 564–577.
- [6] J. Cao, D.-W. Niu and F. Qi, *Convexities of some functions involving the polygamma functions*, Appl. Math. E-Notes **8** (2008), 53–57.
- [7] P. Gao, *A subadditive property of the digamma function*, RGMIA Res. Rep. Coll. **8** (2005), no. 3, Art. 9; Available online at <http://www.staff.vu.edu.au/rgmia/v8n3.asp>.
- [8] B.-N. Guo, R.-J. Chen, and F. Qi, *A class of completely monotonic functions involving the polygamma functions*, J. Math. Anal. Approx. Theory **1** (2006), no. 2, 124–134.
- [9] F. Qi, S. Guo and B.-N. Guo, *Note on a class of completely monotonic functions involving the polygamma functions*, RGMIA Res. Rep. Coll. **10** (2007), no. 1, Art. 5; Available online at <http://www.staff.vu.edu.au/rgmia/v10n1.asp>.

(F. Qi) RESEARCH INSTITUTE OF MATHEMATICAL INEQUALITY THEORY, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

*E-mail address:* [qifeng618@gmail.com](mailto:qifeng618@gmail.com), [qifeng618@hotmail.com](mailto:qifeng618@hotmail.com), [qifeng618@qq.com](mailto:qifeng618@qq.com)

*URL:* <http://qifeng618.spaces.live.com>

(B.-N. Guo) SCHOOL OF MATHEMATICS AND INFORMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO CITY, HENAN PROVINCE, 454010, CHINA

*E-mail address:* [bai.ni.guo@gmail.com](mailto:bai.ni.guo@gmail.com), [bai.ni.guo@hotmail.com](mailto:bai.ni.guo@hotmail.com)

*URL:* <http://guobaini.spaces.live.com>